

*Systems Biology***A knowledge-based approach for predicting gene-disease associations**

Hongyi Zhou and Jeffrey Skolnick*

School of Biology, Georgia Institute of Technology, 950 Atlantic Drive, NW, Atlanta, GA 30332, USA

*To whom correspondence should be addressed.

Associate Editor: Dr. Jonathan Wren

Abstract

Motivation: Recent advances of next-generation sequence technologies have made it possible to rapidly and inexpensively identify gene variations. Knowing the disease association of these gene variations is important for early intervention to treat deadly diseases and provide possible targets to cure these diseases. Genome-wide association studies have identified many individual genes associated with common diseases. To exploit the large amount of data obtained from GWAS studies and leverage our understanding of common as well as rare diseases, we have developed a knowledge-based approach to predict gene-disease associations. We first derive gene-gene mutual information by utilizing the co-occurrence of genes in known gene-disease association data. Subsequently, the mutual information is combined with known protein-protein interaction networks by a boosted tree regression method.

Results: The method called Know-GENE is compared to the method of random walking on the heterogeneous network using the same input data. For a set of 960 diseases, using the same training data in testing in three-fold cross validation, the average recall rate within the top ranked 100 genes by Know-GENE is 65.0% compared to 37.9% by the state of the art random walking on heterogeneous network. This significant improvement is mostly due to the inclusion of knowledge-based mutual information.

Availability: Predictions for genes associated with the 960 diseases are available at <http://cssb2.biology.gatech.edu/knowgene>

Contact: skolnick@gatech.edu

1 Introduction

Complex diseases such as Parkinson's Disease are attributed to both genetic and/or environmental causes (Goldman, 2014). Environmental toxins cause disease through their effects on genes (Qi, et al., 2014). Knowing the genes associated with a disease is useful for preventing and curing the disease. It is also fundamentally important for understanding the biological functions of genes. Genome wide association studies (GWAS) posit that multiple, common small-risk variants interact to cause common diseases (Reich and Lander, 2001; TA, et al., 2010). GWAS relies on testing several hundred thousand common genetic variants found throughout the human genome in large-control cohorts. Over the past few years, many gene loci have been implicated as associated with various human diseases by GWAS (www.genome.gov/gwastudies/) as well as linkage studies (Morton, 1955).

Despite the fruitful insights provided by GWAS, heritability estimates have shown that large proportions of genetic risk underlying complex disease have not yet been explained (Manolio, et al., 2009). This is due to lack of the ability to detect "common disease by rare variants" using a GWAS approach. The introduction of next-generation sequencing technologies allows cost-effective sequencing of entire genomes and has led to the discovery of numerous rare variants in the human genome. The disease associations of these rare variants cannot be inferred by GWAS for further experimental verification. Thus, alternative computational methods that predict the association of gene with a given disease have been developed (Köhler, et al., 2008; Li and Patra, 2010; Natarajan and Dhillon, 2014; Qian, et al., 2014; Singh-Blom, et al., 2013; van Driel, et al., 2006; Vanunu, et al., 2010). The general idea of prediction methods is the "guilt by association" principle (Wolfe, et al., 2005) with respect to a set of known genes related to the given disease. The most frequently used types of evidence for inference of association are (Piro

and Di Cunto, 2012): (a) text mining of the biomedical literature; (b) phenotype relationships such as disease-disease similarity; (c) protein-protein interactions; (d) regulatory information; and (e) gene expression information. For example, Van Driel et al. used a text-mining approach to associate genes with human phenotypes found in the Online Mendelian Inheritance in Man (OMIM) database (Hamosh, et al., 2002; van Driel, et al., 2006). Köhler et al. found gene disease associations by using a global network distance measure – a random walk analysis – for the definition of similarities in protein-protein interaction networks (the “interactome”). Encouragingly, they find that this approach significantly outperforms previous methods based on local distance measures in the interactome (Köhler, et al., 2008). A slightly modified approach called “network propagation” that differs from a random walk only in the normalization of the adjacency matrix W representing the interactome (Qian, et al., 2014; Vanunu, et al., 2010) has also been developed. The random walk method normalizes the adjacency matrix W by columns (the summation of each column equals to one), whereas the network propagation normalizes W by the diagonal matrix: $W_{ij} = W_{ij} / \sqrt{D(i,i)D(j,j)}$, where $D(i,k) = \sum_k W_{ik}$. An extension of the random walk approach is walking on a heterogeneous network that includes protein-protein interaction, disease-disease and gene-disease networks (Li and Patra, 2010). Singh-Blom et al. have developed a truncated version of the random walking on a heterogeneous network by using a limited steps for the walks, but it also includes phenotypes from multiple species (Singh-Blom, et al., 2013). The method uses simple dampening coefficients for longer walks and learns the coefficients for longer walks using a Support Vector Machine (SVM) (Cortes and Vapnik, 1995). Natarajan & Dhillon developed an inductive method that uses a machine learning approach to incorporate different biological sources of evidence such as microarray expression data, gene functional interaction data and disease-related textual data from human as well as other species (Natarajan and Dhillon, 2014). The best performing of the aforementioned methods for prioritizing genes associated with a given disease is the inductive matrix completion developed by Natarajan & Dhillon (Natarajan and Dhillon, 2014). It has an average recall rate of 25% within the top 100 ranked genes. There are also many disease specific methods that focus on a single or group of diseases to prioritize genes for further experimental validation. For a survey of methods for predicting gene-disease association, please see Piro and Di Cunto (Piro and Di Cunto, 2012).

Here, we develop a new type of approach for prioritizing candidate genes associated with a given disease. Our approach applies the idea of word association in the context of texts (Church and Hanks, 1990) to gene-gene association in the context of diseases and then employs gene-gene association to infer gene-disease association. This is a knowledge-based approach that learns gene-gene association propensity in diseases from known gene-disease association. It is also analogous to methods of knowledge-based statistical potentials for protein structure prediction that learn residue-residue or atom-atom pair-wise interaction potentials from experimental protein structures (Lu and Skolnick, 2001; Zhou and Zhou, 2002). Mutual information (Fano, 1961) is used to measure the strength of gene-gene association in a given disease. Owing to the increasing amount of data for known gene-disease associations, the mutual information of gene-gene pairs can be derived from these known associations. Subsequently, mutual information is combined with the properties of the protein-protein physical interaction network by means of boosted tree regression (Roe, et al., 2006). The resulting method called Know-GENE is then benchmarked in three-fold cross-validation (training on 2/3 of the known gene-disease associations and testing on the remaining 1/3) and compared to network propagation, random walking on the interactome as well as on the heterogeneous network based methods. For

960 diseases defined by the Medical Subject Headings (MeSH) ontology with at least 2 known seed genes associated with a given disease (as provided by the OMIM and GWAS databases (Mottaz, et al., 2008; Ramos, et al., 2014; Zhang, et al., 2010)), using the same input evidence (known gene-disease association & interactome), Know-GENE achieves a significantly better recall rate (65.0%) within the top 100 ranked of 15,948 total screened genes compared to the network propagation (19.2%), the random walk on interactome (18.1%), and the random walk on heterogeneous network (37.9%) methods. Thus, Know-GENE is a promising method for prioritizing candidate genes associated with a given disease. Then, we apply Know-GENE in prediction mode (training on all known gene-disease associations) to predict and rank genes in the human exome for each of the 960 diseases. Likewise, our predictions can rank diseases for a given gene, a useful feature for diagnosing diseases in a given mutated gene. It is also useful for predicting and understanding possible side effects of drug targets. Predictions are available for academic users at <http://cssb2.biology.gatech.edu/knowgene>.

2 Methods

This work is based on the assumption that gene-disease association can be inferred from gene-gene functional interactions and protein-protein physical interactions given a set of genes known to be associated with a given disease. We will utilize the known protein-protein physical interaction network and known gene-disease associations as provided by GWAS (Reich and Lander, 2001; TA, et al., 2010) and linkage studies (Morton, 1955).

2.1 Data sets and sources

Protein-protein physical interactions (interactome) are compiled from the HIPPIE database (http://cbdm.mdc-berlin.de/tools/hippie/hippie_current.txt) (Schaefer, et al., 2012) and the work of Menche et al. (<http://www.sciencemag.org/content/347/6224/1257601/suppl/DC1>) (Menche, et al., 2015). In total, there are 246,502 human protein-protein interactions involving 15,948 genes/proteins that are experimentally documented to involve regulatory, metabolic pathway and kinase-substrate interactions. Among these, 3,179 genes are known to be associated with at least one disease. The interactome data can be found at <http://cssb2.biology.gatech.edu/knowgene>.

We then compiled 960 diseases defined by the Medical Subject Heading (MeSH) ontology that have at least 2 associated genes (proteins) in the interactome from two sources as in: (a) Ref. (Zhang, et al., 2010) (<http://www.biomedcentral.com/1755-8794/3/1>) that also uses the MeSH disease definition; and (b) Ref. (Menche, et al., 2015) (<http://www.sciencemag.org/content/347/6224/1257601/suppl/DC1>). The primary sources of these data are from the Genetic Association Databases (GAD) (Becker, et al., 2004), the OMIM database (Hamosh, et al., 2002) and the GWAS databases (Mottaz, et al., 2008; Ramos, et al., 2014). This provides 31,993 gene-disease associations for 3,171 genes. Of these, 283 genes are known to be associated with only one disease, and the disease association of the remaining 15,948 genes is unknown. The known association data can also be found at <http://cssb2.biology.gatech.edu/knowgene>.

2.2 Core genes of a disease

It was found that even with an incomplete interactome, genes associated with given disease tend to interact with each other and form connected clusters (Menche, et al., 2015). A connected cluster consists of genes directly connected (that is interact) to one another in interactome space.

Genes in the largest interacting cluster of a disease are called core genes. To measure if a cluster is formed by chance or due to intrinsic properties of the disease, we conduct the same statistical test as in Ref.(Menche, et al., 2015). For the size S (number of genes) of the cluster, we calculate the z-score:

$$z - score = \frac{S - \langle S^{rand} \rangle}{\sigma(S^{rand})} \quad (1)$$

where $\langle S^{rand} \rangle$ and $\sigma(S^{rand})$ are the average largest cluster size value and standard deviation of randomly picked N_g number of genes in the interaction network. Here, N_g is the number of known genes of the considered disease. To calculate the z-score, we simulate the random process 10,000 times. Each time, N_g genes were randomly selected from the 15,948 screened genes. Then, the size S^{rand} of the largest connected cluster from these N_g genes was obtained. The process was repeated 10,000 times. A cluster with z-score > 1.65 (p -value < 0.05) is considered significant. This leads to 537 of the 960 diseases having a statistically significant cluster. The largest significant cluster of each disease gives the core set of genes: i.e. core genes of the disease. The average number of known genes for the 537 diseases having core genes is 25.5; whereas for the 423 diseases not having core genes, it is 9.4. Thus, a disease having core genes usually has more known genes than diseases lacking core genes.

2.3 Gene-gene pairwise mutual information

Borrowing the idea of measuring word association strength(Church and Hanks, 1990), to measure the strength of functional association of two genes in diseases that might include the effect of direct physical interactions, we use their mutual information (Fano, 1961) defined as:

$$I(g_x, g_y) = \log \frac{P(g_x, g_y)}{P(g_x)P(g_y)} \quad (2)$$

where $P(g_x)$, $P(g_y)$ are the probabilities of observing genes g_x and g_y independently in a given disease, and $P(g_x, g_y)$ is the probability of observing genes g_x and g_y together in a given disease. If there is a genuine association between genes g_x and g_y , then the joint probability $P(g_x, g_y)$ will be larger than that by chance, $P(g_x)P(g_y)$. Thus, $I(g_x, g_y)$ will be > 0 .

In Know-GENE (Knowledge-based approach for predicting gene-disease association), the probabilities $P(g_x)$, $P(g_y)$ are estimated by counting the number of genes g_x , g_y associated with diseases $N(g_x)$ and $N(g_y)$, normalized by the number of diseases, N_d (here, $N_d=960$), and $P(g_x, g_y)$ is estimated by counting the number of co-occurrences of genes g_x , g_y associated with the disease $N(g_x, g_y)$ normalized by N_d .

2.4 Gene-disease association measures

We first consider the network distance of a gene to a given disease. A unit network distance is defined as a path from one protein to another with a direct connection in the interactome. We bin the shortest distances of an unknown gene to all the known genes of a given disease from 1 to 10 and fill each bin with the number of genes known to be associated with the given disease, i.e., we have a vector $(n_1, n_2, \dots, n_{10})$ with $\sum n_i = N_g$, N_g is total number of known genes and n_i is the number of genes in the known set with shortest distance i to the unknown gene. A similar histogram is also done for the core genes (if there is no core gene, all 10 histogram values are zero).

We then consider the functional association strength of an unknown gene g_x to a given disease D that is defined as:

$$S(D, g_x) = \sum_{g_y \in D} P(g_x, g_y) I(g_x, g_y) \quad (3)$$

where the summation is over all known genes of the disease. $P(g_x, g_y)$ and $I(g_x, g_y)$ are the probabilities of observing two genes associated with a disease together and the resulting mutual information (see Eq.(2)), respectively.

To combine the above network distances and functional association strength, we employ the boosted tree regression machine learning method. Boosted tree regression has been employed in many applications(Friedman and Meulman, 2003; Roe, et al., 2006) and has been shown to be much better than support vector machines (SVM)(Cortes and Vapnik, 1995) and random forests (RF)(Breiman, 2001) in predicting genomic breeding values(Ogut, et al., 2011). It involves generating a sequence of decision trees; each grows on the basis of the residuals of all previous trees(Roe, et al., 2006; Thusberg, et al., 2011). Here, a decision tree regression is implemented with a maximal depth of eight. The scoring function is represented as a boosted decision tree(Roe, et al., 2006):

$$f(x) = \sum_{m=1}^{N_{tree}} \epsilon T_m(x) \quad (4)$$

where T_m is a decision tree, ϵ is the shrinkage factor or learning rate, N_{tree} is the number of trees, and x represents a set of features. In this application, N_{tree} is set to 500 and $\epsilon=0.01$. The following 23 features are derived from the above network distances and mutual information:

- 11 features: histogram of network distances to all known genes $(n_1, n_2, \dots, n_{10})$ plus the mean value $\sum (n_i \times i) / N_g$;
- 11 features: histogram of network distances to all core genes $(n_1^c, n_2^c, \dots, n_{10}^c)$ plus the mean value $\sum (n_i^c \times i) / \sum n_i^c$; these values are set to zero for diseases without core genes;
- 1 feature for the functional association strength $S(D, g_x)$ by Eq. (3).

2.5 Training and testing

We considered 960 diseases and 15,948 genes having protein-protein interactions. There are 15,310,080 gene-disease pairs with 31,993 known associations. To test our method, we perform three-fold cross-validation. All gene-disease pairs are randomly partitioned into three approximately equal size sets. For each set, we use the other two sets for training the boosted tree regression model and predict scores for the third, testing set. In the training process, all known associations are assigned an objective function value of 1 and unknown pairs assigned 0. This might introduce a few false negatives because unknown ones could be true associations. However, since the number of unknown associations is much larger than possible true associations, the chance of a false negative is small. In the 2/3 of training data pairs, only 0.2% of total gene-disease pairs are positives (assigned a value of 1). If all pairs are used for training, the model will be overwhelmingly dominated by negative samples. Thus, we shall use only a partial set (~4%) of the negative pairs in the training data for actual training. This results in a ratio of 1:20 positives versus negatives in the actual training data. In either training or testing, when calculating the network distances, the gene is removed from known set of genes for the given disease if it is a known gene of the disease. Any known association is not counted in mutual information derivation by Eq.(2) if it is being tested. The predicted scores are then used to rank genes for each disease. Similarly, one can also use the score to rank diseases of a given gene.

2.6 Comparison to other methods

We shall compare Know-GENE to methods that utilize the same input data (interactome and known set of gene-disease associations) to tease out the effects of different methods. We will not compare methods that use multiple other sources of data (Natarajan and Dhillon, 2014; Singh-Blom, et al., 2013) since they used more data, and the effects of the methods themselves are unclear. We implemented three other methods as described in the corresponding references for comparison: *random walk* (Köhler, et al., 2008), *network propagation* (Vanunu, et al., 2010) and *random walk on a heterogeneous network* (Li and Patra, 2010). *Random walk on a heterogeneous network* requires a disease-disease similarity matrix calculated using MimMiner (van Driel, et al., 2006). In order not to introduce additional data, we used the overlap score to measure disease-disease similarity between disease d_1 and d_2 without using additional input data (Goh, et al., 2007):

$$O(d_1, d_2) = \frac{\text{Number of overlapped genes}}{\sqrt{(\text{number of genes in } d_1)(\text{number of genes in } d_2)}} \quad (5)$$

2.7 Evaluation

For each of the 960 diseases, 15,948 genes are ranked according to their tree regression scores. We evaluated the performance of the methods using the following three criteria: AUC (area under the ROC curve—true positive rate/false positive rate curve), AUPR (area under the precision/recall curve) and recall rate (the ratio (predicted true positives)/(total true positives)) within the top 100 ranked genes. AUC measures discrimination, that is, the ability of the test to correctly classify those with and without the association to the given disease. Of the 15,948 genes, the majority are true negatives (TN) for a given disease, a large number change in the number of false positives (FP) can lead to a small change in the false positive rate (FP/(FP+TN)) used in the ROC analysis (Davis and Goadrich, 2006). On the other hand, AUPR, by comparing false positives to true positives (TP) rather than true negatives, captures the effect of the large number of negative examples on the algorithm's performance (precision=TP/(TP+FP)). Thus, we include AUPR as an additional performance measure. AUPR is sensitive to false positives among the top ranked genes (the region where false positive rate has the smallest value and precision has the largest value). The recall rate gives the experimentalist's expectation of true positives. Based on our earlier work for ligand virtual screening for the human proteome, the recall rate does not depend on whether the known set of genes for a given disease is complete or partial whereas precision, AUC & AUPR will (Zhou, et al., 2015). Thus, recall rate can be considered as having absolute meaning whereas precision, AUC and AUPR are meaningful only for comparing different methods (and are not good for assessing the absolute performance of a method).

3 Results

3.1 Overall performance

The three-fold cross-validation for the 960 diseases is shown in Table 1 in comparison with the other three methods. Know-GENE with AUC=0.967, AUPR=0.405 and recall rate 65.0% is the best for all three measures. Random walk and network propagation methods have very similar results whereas method by walking on the heterogeneous network has much better performance than both the random walk and network propagation. This performance boost is due to the inclusion of information from the gene-disease network. The AUPR and recall rate within the top 100 genes by Know -GENE are almost twice those by walking on a heterogeneous network. We also see that network propagation is slightly better than a random walk. A scatter plot of recall rates

for 960 diseases by Know-GENE versus method of walking on a heterogeneous network is given in Figure 1. Know-GENE wins for 651 diseases whereas walking on the heterogeneous network wins for 129 diseases.

Table 1. Performance of methods for 960 diseases against 15,948 genes

Method	Average AUC	Average AUPR	Average recall rate within the top 100 genes
Random walk	0.780	0.053	18.1%
Network propagation	0.790	0.055	19.2%
Random walk on heterogeneous network	0.930	0.138	37.9%
Know-GENE	0.967	0.405	65.0%
Know-GENE without using mutual information	0.700	0.167	24.0%
Know-GENE without using core gene	0.967	0.352	63.8%

The histograms of recall rates by those two methods are given in Figure 2. For the 960 diseases, Know-GENE has 71.4% of diseases with a recall rate $\geq 50\%$, whereas walking on heterogeneous network has 34.3% of diseases with a recall rate $\geq 50\%$. The p -value for the difference between recall rates of Know-GENE and those of walking on heterogeneous network is 2.8×10^{-92} . Thus, Know-GENE gives significant improvement over extant methods using exactly the same input data.




Fig. 1. Scatter plot of recall rates within the top 100 genes by Know-GENE versus the walking on heterogeneous network approach.




Fig. 2. Fractions of diseases having recall rates within the top 100 genes \geq given value by Know-GENE and the walking on heterogeneous network approach, respectively.

3.2 Effects of mutual information and core genes

We next examine the effects of some factors in Know-GENE by removing from the features: (a) the mutual information term or (b) the network distances to core genes. The results are also given in Table 1, and clearly indicate that without the mutual information term, Know-GENE performs significantly worse. Thus, the mutual information term derived from the gene-disease network contributes the most to the good performance of Know-GENE. This is consistent with the method by walking on the heterogeneous network that gets boosted performance by including the gene-disease network. The fact that Know-GENE performs significantly better than walking on the heterogeneous network indicates that mutual information captures more important information than the random walk approach. The terms related to core genes slightly affect the AUPR (from 0.405 to 0.352) and recall rate (from 65.0% to 63.8%).

In Table 2, we separately examined the performance of the methods for the 537 diseases having core genes and 423 diseases with no core genes. All methods have better performance for diseases having core genes. The performance differences between the two data sets are relatively larger for random walk and network propagation methods where known gene-disease associations are not utilized. Know-GENE has the best performance for all criteria for both data sets. It has a 70.7% average recall rate for the 537 diseases having core genes.

Table 2. Performance of methods to predict disease association of the 15,948 genes for diseases with or without core genes

Method	Average AUC	Average AUPR	Average recall rate within the top 100 genes
537 diseases having core genes			
Random walk	0.810	0.082	25.7%
Network propagation	0.823	0.086	27.1%
Random walk on heterogeneous network	0.957	0.179	41.2%
Know-GENE	0.980	0.533	70.7%
423 diseases not having core genes			
Random walk	0.741	0.016	8.4%
Network propagation	0.748	0.015	9.0%
Random walk on heterogeneous network	0.896	0.086	33.7%
Know-GENE	0.951	0.243	57.7%

3.3 Test on singleton genes

Next, we analyze the performance of methods for *singleton* genes (Natarajan and Dhillon, 2014; Singh-Blom, et al., 2013) defined as having only one known disease association in the data. In this case, the gene will have no mutual information contribution at training and testing times because it is removed from the known association with a given disease. Therefore, the test on *singleton* genes tests the ability of the method to predict associations for genes that have no known association to any disease. The test results are compiled in Table 3 using measures of the average rank of genes and number of genes ranked within the top 100 of 15,948 genes for a given disease. Table 3 shows that all methods perform much poorer (the best recall rate by network propagation method is 32/283=11%) for *singleton* genes than that for genes known to be associated with diseases. Especially for Know-GENE and walking on heterogeneous network that utilized known gene-disease associations, the recall rate is around 1%. This is because these methods favor genes that have known gene-disease associations and thus drag down the rela-

tive ranks of those genes have no known gene-disease associations. Therefore, in practice, if a gene has no known disease association, one should use methods like network propagation, random walk, or Know-GENE without mutual information whose recall rate is 26/283 or 9%

Table 3. Performance of different methods to assign diseases for 283 *singleton* genes

Method	Average rank	# of genes ranked within the top 100 genes
Random walk	5485	24
Network propagation	5014	32
Random walk on heterogeneous network	6948	3
Know-GENE	8005	3
Know-GENE without using mutual information	5699	26
Know-GENE without using core gene	7969	1

3.4 Predicting new genes associated with a given disease

In the above benchmark tests, we examine only the top 100 ranked genes for a given disease. In practice, some of the diseases are associated with fewer than 100 genes and some diseases might have associations with more than 100 genes. We thus optimize a cutoff score that will give the best binary classification (associated/not associated) measured by the Matthew's Correlation Coefficient (MCC) for the training data. This results in a cutoff score value of 0.45 and a MCC of 0.878 for the training data. We then use all the 15,278,087 predicted gene-disease pair scores with unknown associations as a random score distribution and fit the distribution to an Extreme Value Distribution with $\mu=0.002$ and $\sigma=0.0122$. A cutoff of 0.45 corresponds to a p -value of 2.2×10^{-16} . With a 0.45 cutoff, Know-GENE will have an average per disease recall of 53.3%, slightly worse than 65.0% within top 100 ranked genes (notice that AUC & AUPR are cutoff independent).

Now, we examine predictions of new genes for specific diseases. One example is Parkinson's Disease (PD) (Goldman, 2014). In our test data, there are 38 genes known to be associated with PD. Using a cutoff score of 0.45, Know-GENE predicts 149 genes associated with PD. 32 (84.2%) of the 38 known associated genes are among the predictions. For the top 10 predicted genes with unknown PD association, we searched the literature to support or disapprove our predictions, with the results compiled in Table 4. Eight (80%) of the top 10 genes have supporting evidence giving a prediction precision of ~80% for this particular disease.

Table 4. Predicted top 10 new genes for Parkinson's Disease by Know-GENE

Gene	Score	Evidence of support or disapprove
NUP62_HUMAN	0.971	Supported by ref.(Zatloukal, et al., 2002)
HD_HUMAN	0.953	Causes Huntington's disease that shares the same malfunctions within the motor sector of the nervous system as PD (Delcomyn, 1998); http://serendip.brynmawr.edu/bb/neuro/neuro98/202s98-paper3/Sangaramoorthy3.html

PANK2_HUMAN	0.953	Disapproved by ref.(Klopstock, et al., 2005)
TOR1A_HUMAN	0.952	Supported by ref.(Leung, et al., 2001)
TOM40_HUMAN	0.935	Supported by ref.(Bender, et al., 2013)
STX6_HUMAN	0.922	Disapprove by ref.(Trinh, et al., 2013)
THAP1_HUMAN	0.890	Support: THAP1 gene is part of a family of THAP proteins that bind specific DNA sequences and regulate cell proliferation through the pRB/E2F cell cycle target genes, a pathway recently proposed to be involved in cell death in Parkinson disease(Höglinger, et al., 2007; Houlden, et al., 2010).
IRF4_HUMAN	0.875	Supported by ref.(Soreq, et al., 2008)
E2AK3_HUMAN	0.875	Supported by ref.(Dzamko, et al., 2014)
A4_HUMAN	0.862	Supported by ref.(Schulte, et al., 2015)

3.5 Predicting new diseases associated with a given gene

We can also examine the prediction of new diseases associated with a given gene. Diseases having a score ≥ 0.45 are used as predictions for a given gene. First, we examine the overall performance of disease prediction for a given gene by ranking diseases. The average AUC, AUPR and recall rate for 2,888 genes having at least 2 known associated diseases are 0.969, 0.644 and 77.1%, respectively. We then examine the overall frequency of a gene to be associated with diseases and check its correlation with the number of interactions the corresponding protein has. The number of interactions of a protein can be derived from the interactome. Figure 3 shows the average number of interactions a gene has versus the number of diseases associated with a given gene \geq given threshold. The Pearson's Correlation Coefficient (C.C.) between the average number of interactions and threshold number of diseases when using known associations is 0.906, and when using predicted associations by Know-GENE with the cutoff of 0.45, it is 0.994, respectively. On average, proteins having more physical interactions tend to be associated with more diseases.




Fig. 3. Average number of interactions a gene has versus the number of diseases associated with the gene \geq given value. The Pearson's Correlation Coefficient between the average number of interactions and threshold number of diseases when using known gene-disease associations is

0.906; when using predicted associations by Know-GENE with a score ≥ 0.45 , it is 0.994, respectively.

One example of predicted new diseases for a given gene is for p53, a tumor suppressor gene. In our data, p53 is known to be associated with 100 diseases, and 84 of them have a score ≥ 0.45 . Another 135 diseases also have a score ≥ 0.45 for p53, but these have unknown associations with p53. We list the top 10 new diseases predicted for the p53 gene in Table 5 along with supporting evidence. Nine diseases are supported, and one disapproved by literature.

Table 5. Predicted top 10 new diseases for the p53 gene by Know-GENE

Disease	Score	Evidence of support or disapprove
Female urogenital diseases and pregnancy complications	0.986	Supported by ref.(Fraga, et al., 2014)
Intestinal Diseases	0.981	Supported by ref.(Hussain, et al., 2000)
Congenital Abnormalities	0.967	Supported by ref.(Barkić, et al., 2009)
Skin and connective tissue diseases	0.965	Supported by ref.(Menke, et al., 2000)
Neoplasms, nerve tissue	0.961	Supported by ref.(Barbareschi, et al., 1992)
Skin Diseases	0.960	Supported by ref.(Batinac, et al., 2004)
Neoplasms, germ cell and embryonal	0.947	Supported by ref.(Lewis, et al., 1994)
Metabolism, Inborn Errors	0.945	Supported by ref.(Wilkins, et al., 2013)
Neuroectodermal tumors	0.944	Disapprove by ref.(Raffel, et al., 1993)
Male Urogenital Diseases	0.941	Supported by ref.(Castrén, et al., 1998)

We also find that disease recall rate is positively correlated with the number of known disease associations a given gene has. The Pearson's Correlation Coefficient is 0.173 for 2,888 genes, which corresponds to a p -value of 7.7×10^{-21} . For the 283 singleton genes, the average AUC, AUPR and recall rate by Know-GENE are 0.431, 0.012 and 0.0%, respectively, while the average AUC, AUPR and recall rate by Know-GENE without using mutual information are 0.841, 0.050 and 24.7%, respectively (using the cutoff=0.32 obtained by optimizing MCC of the training set). Thus, Know-GENE is not good for predicting disease association of genes not seen in any of the diseases used for deriving the mutual information. In such cases, it would be better to apply Know-GENE without using mutual information or network propagation approach or the approach of simply walking on interactome instead of heterogeneous network.

Finally, Using Know-GENE in prediction mode, i.e. training on all data and making predictions for all genes, we provide all prediction results and a stand-alone program for academic users at <http://cssb2.biology.gatech.edu/knowgene>. The stand-alone program can be used to plug in new diseases with known genes for re-training the model and for predicting new genes associated with the already included diseases. Instructions as to how to use the program are also provided.

4 Discussion

We have developed a knowledge-based approach Know-GENE for prioritizing genes associated with given disease. Likewise, it can also prioritize diseases associated with a given gene. The novelty of Know-GENE is in the derivation of gene-gene mutual information from the co-occurrence frequency of pairs of genes in a large number of diseases with a known set of gene-disease associations. With this novel technique, Know-GENE performs much better than the best existing method of walking on the heterogeneous network using exactly the same input information (recall rate within top 100 ranked genes: 65.0% vs. 37.9%). Both Know-GENE and walking on the heterogeneous network utilize information from the known gene-disease associations and perform much better than methods without using this information. For example, the network propagation method has a recall rate of 19.2%, which is about half that by the method of random walk on the heterogeneous network. All tested methods perform better for diseases having core genes than for those lacking core genes. When used for prioritizing diseases for a given gene, Know-GENE has an average recall rate of 77.1%.

A disadvantage of Know-GENE as well as the method of walking on the heterogeneous network is that it does not perform well for genes that are not present in any of the diseases used in deriving the mutual information (see Table 3 test on singletons). In practice, for such genes, we can choose an alternative method such as Know-GENE without using mutual information, or the network propagation method.

Possible further improvement of Know-GENE could come from enriching more genes with more verified disease associations and including them in deriving mutual information. These could be fulfilled by literature searches for associations with high Know-GENE prediction scores and then using the supported associations in training. This process could be iterated. For those genes without any known disease associations, other sources for deriving their mutual information are sought. For example, one possibility is the co-variation of gene expression data in individuals with a given disease. Co-occurrence of gene pairs in pathways is another possible source. These and other alternatives will be explored in future work.

Acknowledgements

The authors thank Dr. Bartosz Ilkowski for managing the cluster on which this work was conducted.

Funding

This work has been supported by the National Institutes of Health grant No GM-48835.

Conflict of Interest: none declared.

References

Barbareschi, M., *et al.* p53 protein expression in central nervous system neoplasms. *J Clin Pathol.* 1992;45(7):583-586.

Barkić, M., *et al.* The p53 Tumor Suppressor Causes Congenital Malformations in Rpl24-Deficient Mice and Promotes Their Survival. *Mol. Cell. Biol.* 2009;29(10):2489-2504.

Batinac, T., *et al.* p53 protein expression and cell proliferation in non-neoplastic and neoplastic proliferative skin diseases. *Tumori.* 2004;90(120-7).

Becker, K., *et al.* The genetic association database. *Nat Genet* 2004;36(5):431-432.

Bender, A., *et al.* TOM40 Mediates Mitochondrial Dysfunction Induced by α -Synuclein Accumulation in Parkinson's Disease. *PLoS One* 2013;8(4):e62277.

Breiman, L. Random Forests. *Machine Learning* 2001;45:5--32.

Castrén, K., *et al.* Absence of p53 mutations in benign and pre-malignant male genital lesions with over-expressed p53 protein. *Int J Cancer* 1998;77(5):674-678.

Church, K.W. and Hanks, P. Word association norms, mutual information, and lexicography. *Comput. Linguist.* 1990;16(1):22-29.

Cortes, C. and Vapnik, V. Support-vector networks. *Machine Learning* 1995;20(3).

Davis, J. and Goadrich, M. The relationship between Precision-Recall and ROC curves. *Proceedings of the 23rd international conference on Machine learning. ACM* 2006:233-240.

Delcomyn, F. Foundations of Neurobiology. In, *Foundations of Neurobiology.* New York: W.H. Freeman and Company; 1998. p. 436-437.

Dzamko, N., *et al.* Parkinson's disease-implicated kinases in the brain; insights into disease pathogenesis. *Front Mol Neurosci.* 2014;7:57.

Fano, R. Transmission of Information: A Statistical Theory of Communications. Cambridge, MA: MIT Press; 1961.

Fraga, L., *et al.* p53 signaling pathway polymorphisms associated to recurrent pregnancy loss. *Mol. Biol. Rep.* 2014;41(3):1871-1877.

Friedman, J.H. and Meulman, J.J. Multiple additive regression trees with application in epidemiology. *Statistics in Medicine* 2003;22:1365--1383.

Goh, K.-I., *et al.* The human disease network. *PNAS* 2007;104(21):8685-8690.

Goldman, S.M. Environmental Toxins and Parkinson's Disease. *Annu. Rev. Pharmacol. Toxicol.* 2014;54:141-164.

Hamosh, A., *et al.* Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders. *Nucl. Acids Res.* 2002;30:52--55.

Höglinger, G., *et al.* The pRb/E2F cell-cycle pathway mediates cell death in Parkinson's disease. *Proc Natl Acad Sci U S A.* 2007;104(9):3585-3590.

Houlden, H., *et al.* THAP1 mutations (DYT6) are an additional cause of early-onset dystonia. *Neurology* 2010;74(10):846-850.

Hussain, S., *et al.* Increased p53 mutation load in noncancerous colon tissue from ulcerative colitis: a cancer-prone chronic inflammatory disease. *Cancer Res.* 2000;60(13):3333-3337.

Klopstock, T., *et al.* Mutations in the pantothenate kinase gene PANK2 are not associated with Parkinson disease. *Neurosci Lett.* 2005;379(3):195-198.

Köhler, S., *et al.* Walking the interactome for prioritization of candidate disease genes. *Am J Hum Genet.* 2008;82(4):949--958.

Leung, J., *et al.* Novel mutation in the TOR1A (DYT1) gene in atypical early onset dystonia and polymorphisms in dystonia and early onset parkinsonism. *Neurogenetics* 2001;3(3):133-143.

Lewis, D., *et al.* Immunohistochemical expression of P53 tumor suppressor gene protein in adult germ cell testis tumors: clinical correlation in stage I disease. *J Urol.* 1994;152(2 Pt 1):418-423.

Li, Y. and Patra, J. Genome-wide inferring gene-phenotype relationship by walking on the heterogeneous network. *Bioinformatics* 2010;26(9):1219-1224.

Lu, H. and Skolnick, J. A distance-dependent atomic knowledge-based potential for improved protein structure selection. *Proteins* 2001;44:223--232.

Manolio, T., *et al.* Finding the missing heritability of complex diseases. *Nature* 2009;461:747-753.

Menche, J., *et al.* Uncovering disease-disease relationships through the incomplete interactome. *Science* 2015;347:841.

Menke, D., *et al.* Lymphomas in patients with connective tissue disease. Comparison of p53 protein expression and latent EBV infection in patients immunosuppressed and not immunosuppressed with methotrexate. *Am J Clin Pathol.* 2000;113(3):212-218.

Morton, N. Sequential tests for the detection of linkage. *Am J Hum Genet.* 1955;7(3):277-318.

Mottaz, A., *et al.* Mapping proteins to disease terminologies: from UniProt to MeSH. *BMC Bioinformatics.* 2008;9(Suppl 5):S3.

Natarajan, N. and Dhillon, I.S. Inductive matrix completion for predicting gene-disease associations. *Bioinformatics* 2014;30(12):i60--i68.

Ogut, J., Piepho, H. and Schiulz-Streeck, T. A comparison of random forests, boosting and support vector machines for genomic selection. *BMC Proceedings* 2011;5(Suppl 3):S11.

Piro, R. and Di Cunto, F. Computational approaches to disease-gene prediction: rationale, classification and successes. *FEBS J.* 2012;279(5):678-696.

Qi, Z., Miller, G.W. and Voit, E.O. Rotenone and paraquat perturb dopamine metabolism: a computational analysis of pesticide toxicity. *Toxicology* 2014;315:92-101.

Qian, Y., *et al.* Identifying disease associated genes by network propagation. *BMC Systems Biology* 2014;8(suppl 1):S6.

Raffel, C., *et al.* Absence of p53 mutations in childhood central nervous system primitive neuroectodermal tumors. *Neurosurgery* 1993;33(2):301-305.

-
- Ramos, E., *et al.* Phenotype-Genotype Integrator (PheGenI): synthesizing genome-wide association study (GWAS) data with existing genomic resources. *Eur J Hum Genet.* 2014;22(1):144-147.
- Reich, D. and Lander, E. On the allelic spectrum of human disease. *Trends Genet.* 2001;17(9):502-510.
- Roe, B.P., Yang, H.-J. and Zhu, J. Boosted Decision Trees, A Powerful Event Classifier. In, *Statistical Problems in Particle Physics, Astrophysics and Cosmology.* . 2006. p. 139.
- Schaefer, M.H., *et al.* HIPPIE: Integrating Protein Interaction Networks with Experiment Based Quality Scores. *Plos One* 2012;7(2):e31826.
- Schulte, E., *et al.* Rare variants in β -Amyloid precursor protein (APP) and Parkinson's disease. *Eur J Hum Genet.* 2015;23(10):1328-1333.
- Singh-Blom, U.M., *et al.* Prediction and Validation of Gene-Disease Associations Using Methods Inspired by Social Network Analyses. *PLOS ONE* 2013;8(5):e58977.
- Soreq, L., *et al.* Advanced microarray analysis highlights modified neuro-immune signaling in nucleated blood cells from Parkinson's disease patients. *Journal of Neuroimmunology* 2008;201:227-236.
- TA, M., Guttmacher, A.E. and Manolio, T.A. Genomewide association studies and assessment of the risk of disease. *N. Engl. J. Med.* 2010;363(2):166-176.
- Thusberg, J., Olatubosun, A. and Vihinen, M. Performance of mutation pathogenicity prediction methods on missense variants. *Hum Mutat.* 2011;32(4):356--368.
- Trinh, J., *et al.* STX6 rs1411478 is not associated with increased risk of Parkinson's disease. *Parkinsonism Relat Disord.* 2013;19(5):563-565.
- van Driel, M., *et al.* A text-mining analysis of the human phenome. *Eur J Hum Genet.* 2006;14(5):535-542.
- Vanunu, O., *et al.* Associating Genes and Protein Complexes with Disease via Network Propagation. *PLOS Computational Biology* 2010;6(1):e1000641.
- Wilkins, B.J., *et al.* p53-Mediated Biliary Defects Caused by Knockdown of cirh1a, the Zebrafish Homolog of the Gene Responsible for North American Indian Childhood Cirrhosis. *PLOS ONE* 2013;8(10): e77670.
- Wolfe, C., Kohane, I. and Butte, A. Systematic survey reveals general applicability of "guilt-by-association" within gene coexpression networks. *BMC Bioinformatics.* 2005;6:227.
- Zatloukal, K., *et al.* p62 Is a common component of cytoplasmic inclusions in protein aggregation diseases. *Am J Pathol.* 2002;160(1):255-263.
- Zhang, Y., *et al.* Systematic analysis, comparison, and integration of disease based human genetic association data and mouse genetic phenotypic information. *BMC Medical Genomics* 2010;3:1.
- Zhou, H., Gao, M. and Skolnick, J. Comprehensive prediction of drug-protein interactions and side effects for the human proteome. *Scientific Reports* 2015;5:11090.
- Zhou, H. and Zhou, Y. Distance-scaled, finite ideal-gas reference state improves structure-derived potentials of mean force for structure selection and stability prediction. *Protein Science* 2002;11 2714--2726.